Sonata generic

Medically reviewed by Sophia Entringer, PharmD. Last updated on April 21, Sonata zaleplon is a sedative, also called a hypnotic. It affects chemicals in your brain that may be unbalanced in people with sleep problems insomnia. Sonata is used to treat insomnia. Sonata may cause a severe allergic reaction. Stop taking this medicine and get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Some people using Sonata have engaged in activity such as driving, eating, or making phone calls and later having no memory of the activity. Serious injury or death could occur if you walk or drive while you are not fully awake. If this happens to you, stop taking this medication and talk with your doctor about another treatment for your sleep disorder. Sonata can cause side effects that may impair your thinking or reactions. You may still feel sleepy the morning after taking the medication. Do not drink alcohol while you are taking this medication. It can increase some of the side effects of zaleplon, including drowsiness. Zaleplon may be habit-forming. Keep the medication in a place where others cannot get to it. Selling or giving away Sonata is against the law. You may have withdrawal symptoms if you stop taking this medication after taking it over several days in a row. Do not stop taking Sonata suddenly without first talking to your doctor. You may need to use less and less before you stop the medication completely. To make sure this medicine is safe for you, tell your doctor if you have ever been addicted to drugs or alcohol. You should not use Sonata if you are allergic to zaleplon, or if you have ever taken sleep medicine and engaged in activity you later don't remember. The sedative effects of zaleplon may be stronger in older adults. Accidental falls are common in elderly patients who take sedatives. Use caution to avoid falling or accidental injury while you are taking Sonata. Take Sonata exactly as it was prescribed for you. Follow all directions on your prescription label and read all medication guides or instruction sheets. Misuse can cause addiction, overdose, or death. Selling or giving away this medicine is against the law. Sonata will make you fall asleep. Never take this medication during your normal waking hours, unless you have several hours to dedicate to sleeping. Sonata is for short-term use only. Do not take this medicine for longer than 5 weeks without your doctor's advice. Call your doctor if your symptoms do not improve after 7 to 10 days of treatment, or if they get worse. Do not stop using Sonata suddenly after taking it over several days in a row, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine. Store at room temperature away from moisture, heat, and light. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription. Insomnia symptoms may also return after you stop taking Sonata, and may be even worse than before. Tell your doctor if you still have worsened insomnia after the first few nights without taking this medicine. Use: Short-term treatment of insomnia. Since Sonata is taken only at bedtime, you will not be on a frequent dosing schedule. Never take this medication if you do not have several hours to sleep before being active again. Seek emergency medical attention or call the Poison Help line at An overdose of zaleplon can be fatal, especially when it is taken together with other medications that can cause drowsiness. Overdose symptoms may include severe drowsiness, confusion, loss of coordination, severe muscle weakness, weak or shallow breathing, fainting. Avoid taking Sonata during travel, such as to sleep on an airplane. You may be awakened before the effects of the medication have worn off. Amnesia forgetfulness is more common if you do not get the equivalent of a full night's sleep after taking Sonata. Avoid taking Sonata with food or right after eating a high-fat or heavy meal. This will make it harder for your body to absorb the medicine. Zaleplon can cause side effects that may impair your thinking or reactions. Until you know how this medicine will affect you during waking hours, be careful if you drive, operate machinery, pilot an airplane, or do anything that requires you to be awake and alert. Zaleplon may cause a severe allergic reaction. Get emergency medical help if you have signs of an allergic reaction to Sonata : hives ; difficult breathing; nausea and vomiting ; swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, stop taking Sonata and call your doctor right away. This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. Using Sonata with other drugs that make you drowsy or slow your breathing can cause dangerous side effects or death. Ask your doctor before using opioid medication, other sleep medicine, a muscle relaxer, or medicine for anxiety or seizures. Other drugs may interact with zaleplon, including prescription and over-the-counter medicines, vitamins , and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using. Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Sonata only for the indication prescribed. Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Skip to Content. Treatments Insomnia Sonata

Print Share. Sonata dosage more detail. Sonata side effects more detail. Sonata drug interactions more detail. Drug Status Availability Prescription only Rx. Drug Class. Miscellaneous anxiolytics, sedatives and hypnotics. Related Drugs. Subscribe to our newsletters. FDA Safety Alerts for all medications. Daily MedNews. Weekly Drug News Roundup. Monthly Newsletter. I accept the Terms and Privacy Policy. Email address. Select one or more newsletters to continue. Harvard Health Publishing. Related: Insomnia. Mayo Clinic. Sonata is a brand name of zaleplon , approved by the FDA in the following formulation s :. Note: Fraudulent online pharmacies may attempt to sell an illegal generic version of Sonata. These medications may be counterfeit and potentially unsafe. If you purchase medications online, be sure you are buying from a reputable and valid online pharmacy. Ask your health care provider for advice if you are unsure about the online purchase of any medication. Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. More drug approvals. Skip to Content. Generic Drugs Sonata Print Share. Patent and Trademark Office and assigns exclusive legal right to the patent holder to protect the proprietary chemical formulation. The patent assigns exclusive legal right to the inventor or patent holder, and may include entities such as the drug brand name, trademark, product dosage form, ingredient formulation, or manufacturing process A patent usually expires 20 years from the date of filing, but can be variable based on many factors, including development of new formulations of the original chemical, and patent infringement litigation. Drug Exclusivity Exclusivity is the sole marketing rights granted by the FDA to a manufacturer upon the approval of a drug and may run simultaneously with a patent. Exclusivity periods can run from days to seven years depending upon the circumstance of the exclusivity grant. By designating a single reference listed drug as the standard to which all generic versions must be shown to be bioequivalent, FDA hopes to avoid possible significant variations among generic drugs and their brand name counterpart. AB Products meeting necessary bioequivalence requirements. Multisource drug products listed under the same heading i. In certain instances, a number is added to the end of the AB code to make a three character code i. Three-character codes are assigned only in situations when more than one reference listed drug of the same strength has been designated under the same heading. Two or more reference listed drugs are generally selected only when there are at least two potential reference drug products which are not bioequivalent to each other. If a study is submitted that demonstrates bioequivalence to a specific listed drug product, the generic product will be given the same three-character code as the reference listed drug it was compared against. Drug Status Availability Prescription only Rx. Drug Class. Miscellaneous anxiolytics, sedatives and hypnotics. Recently Approved. Ukoniq Ukoniq umbralisib is a dual inhibitor of phosphoinositide 3 kinase PI3K Subscribe to our newsletters. FDA Safety Alerts for all medications. Daily MedNews. Weekly Drug News Roundup. Monthly Newsletter. I accept the Terms and Privacy Policy. Email address. Select one or more newsletters to continue. A drug patent is assigned by the U. Exclusivity is the sole marketing rights granted by the FDA to a manufacturer upon the approval of a drug and may run simultaneously with a patent. A Reference Listed Drug RLD is an approved drug product to which new generic versions are compared to show that they are bioequivalent. Products meeting necessary bioequivalence requirements. Medically reviewed by Drugs. Last updated on Dec 3, The dose of Sonata should be individualized. The recommended dose of Sonata for most nonelderly adults is 10 mg. For certain low weight individuals, 5 mg may be a sufficient dose. Although the risk of certain adverse events associated with the use of Sonata appears to be dose dependent, the 20 mg dose has been shown to be adequately tolerated and may be considered for the occasional patient who does not benefit from a trial of a lower dose. Doses above 20 mg have not been adequately evaluated and are not recommended. Elderly patients and debilitated patients appear to be more sensitive to the effects of hypnotics, and respond to 5 mg of Sonata. The recommended dose for these patients is therefore 5 mg. Doses over 10 mg are not recommended. Hepatic insufficiency: Patients with mild to moderate hepatic impairment should be treated with Sonata 5 mg becau

renault clio immobiliser wiring diagram
2000 toyota 4runner radio
honda gx620 carburetor diagram

se clearance is reduced in this population. Sonata is not recommended for use in patients with severe hepatic impairment. Renal insufficiency: No dose adjustment is necessary in patients with mild to moderate renal impairment. Sonata has not been adequately studied in patients with severe renal impairment. Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Skip to Content. Sonata Dosage Medically reviewed by Drugs. Special Populations Elderly patients and

debilitated patients appear to be more sensitive to the effects of hypnotics, and respond to 5 mg of Sonata. Drug Status Availability Prescription only Rx. Drug Class. Miscellaneous anxiolytics, sedatives and hypnotics. Related Drugs. Subscribe to our newsletters. FDA Safety Alerts for all medications. Daily MedNews. Weekly Drug News Roundup. Monthly Newsletter. I accept the Terms and Privacy Policy. Email address. Select one or more newsletters to continue.